

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/335877398>

Density-dependent growth response and sensitivity of *Meloidogyne incognita* to *Maerua angolensis* and *Tabernaemontana elegans* : Card model

Article in *Acta Agriculturae Scandinavica, Section B - Soil & Plant Science* · September 2019

DOI: 10.1080/09064710.2019.1663913

CITATIONS

0

READS

103

6 authors, including:



Mbokota Candy Khosa

Agricultural Research Council, South Africa

6 PUBLICATIONS 17 CITATIONS

[SEE PROFILE](#)



Zanele Dube

University of KwaZulu-Natal

15 PUBLICATIONS 28 CITATIONS

[SEE PROFILE](#)



Malefa Tselanyane

Council for Scientific and Industrial Research, South Africa

21 PUBLICATIONS 109 CITATIONS

[SEE PROFILE](#)



Gerda Fouche

Council for Scientific and Industrial Research, South Africa

53 PUBLICATIONS 605 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Isolation of compounds [View project](#)



The search for new anti-aging ingredients [View project](#)



Density-dependent growth response and sensitivity of *Meloidogyne incognita* to *Maerua angolensis* and *Tabernaemontana elegans*: Card model

M. C. Khosa, Z. P. Dube, M. Tselanyane, G. Fouche, D. De Waele & M. S. Daneel

To cite this article: M. C. Khosa, Z. P. Dube, M. Tselanyane, G. Fouche, D. De Waele & M. S. Daneel (2019): Density-dependent growth response and sensitivity of *Meloidogyne incognita* to *Maerua angolensis* and *Tabernaemontana elegans*: Card model, Acta Agriculturae Scandinavica, Section B — Soil & Plant Science, DOI: [10.1080/09064710.2019.1663913](https://doi.org/10.1080/09064710.2019.1663913)

To link to this article: <https://doi.org/10.1080/09064710.2019.1663913>



Published online: 17 Sep 2019.



Submit your article to this journal [↗](#)



Article views: 7



View related articles [↗](#)



View Crossmark data [↗](#)

SHORT COMMUNICATION



Density-dependent growth response and sensitivity of *Meloidogyne incognita* to *Maerua angolensis* and *Tabernaemontana elegans*: Card model

M. C. Khosa^a, Z. P. Dube^b, M. Tselanyane^c, G. Fouche^d, D. De Waele^{e,f} and M. S. Daneel^a

^aAgricultural Research Council-Tropical and Subtropical Crops, Nelspruit, South Africa; ^bSchool of Biology and Environmental Sciences, University of Mpumalanga, Mbombela, Republic of South Africa; ^cBiosciences Unit, Council of Scientific and Industrial Research, Pretoria, South Africa; ^dDepartment of Chemistry, University of Pretoria, Pretoria, South Africa; ^eUnit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa; ^fLaboratory of Tropical Crop Improvement, Department of Biosystems, Faculty of Bioscience Engineering, University of Leuven (KU Leuven), Leuven, Belgium

ABSTRACT

Curve-fitting Allelochemical Response Dosage (CARD) computer-based model has proven to be a valuable tool in the definition of dosage response relationships between organisms and various plant extracts. Its use in the explanation of the relationship between *Maerua angolensis* and *Tabernaemontana elegans* concentrations and *Meloidogyne incognita* second-stage juvenile (J2) hatch and mortality has not been reported. The objective of this study was to use the CARD model in the determination of *M. incognita* sensitivity, density-dependent growth (DDG) patterns and minimum inhibition concentrations of *M. angolensis* and *T. elegans* plant extracts on juvenile hatch and mortality. The CARD model quantified concentration ranges of the two plant extracts that could stimulate ($D_m - R_h$), saturate ($R_h - D_0$) and inhibit ($D_0 - D_{100}$), J2 hatch and mortality. *Meloidogyne incognita* J2 hatch was moderately sensitive to *M. angolensis* and *T. elegans* plant extracts with average sensitivity ranking of 20 and 19 units, respectively. J2 mortalities were moderately sensitive to *T. elegans* average sensitivity ranking of 15 units, whereas highly sensitive to *M. angolensis* at 2 units. The minimum juvenile hatch inhibition and lethal concentrations for the two extracts were comparable and very low at 0.07, 0.06 and 0.13, 0.94 mg mL⁻¹, respectively. In conclusion, the low minimum concentrations for *M. incognita* J2 hatch inhibition and lethal concentrations of *M. angolensis* and *T. elegans* plant extracts and moderate sensitivity values provide substantial evidence of high potency of the two extracts in the potential management of *M. incognita*.

ARTICLE HISTORY

Received 12 June 2019
Accepted 26 August 2019

KEYWORDS

Bead-bean; root-knot nematode; medicinal plant; phytonematicides; sensitivity index; toad tree

Introduction

Mathematical modelling has had extensive use in crop protection, especially in the development of integrated pest management strategies (Walczak 2011; Nayak et al. 2018). Most of these models focussed on disease risk assessments (Gutsche and Rossberg 1997) and impact of pests on yield (Herbert et al. 1992; Donatelli et al. 2017) with few developed to predict the impact of synthetic pesticide applications and threshold levels for spraying on the environment and crop being protected (Fisker et al. 2007; Walczak 2011). There are currently limited models developed for quick preliminary evaluation of phytopesticide efficacy (Liu et al. 2003) even though they have been proven to be viable alternatives to synthetic chemicals in pest management (Mashela et al. 2015).

Liu et al. (2003) developed a computer-based model that assessed the response of an organism when exposed to varying concentrations of allelochemicals.

The model has been adapted for use in phytonematicide dosage trials (Mafeo 2012; Pelinganga 2013; Mashela et al. 2015; Dube 2016). The initial adoption of the model was focused on quantifying all stages of the density-dependent growth (DDG) patterns of different crop plants to increasing concentrations of phytonematicides, values which were later used to determine a concentration of a phytonematicide which stimulates plant growth, while suppressing population densities of the target pest (Mafeo 2012; Pelinganga 2013). Dube (2016) demonstrated that the model also has potential in *in vitro* trials of phytonematicides for quick screening through the establishment of toxicity and minimum inhibition concentration.

Curve-fitting Allelochemical Response Dosage (CARD) model is based on mathematical model of curvilinear regression equations. The model uses several generated biological indices to formulate a density-dependent growth response of an organism to allelochemicals and

various other variables (Liu et al. 2003). The two most used biological indices of the model are those that characterise the organism's response towards allelochemicals in a DDG pattern and sensitivity index (k) (Mashela et al. 2015). The density-dependent response patterns are critical in the determination of the optimum and/or minimum inhibition concentration of allelochemicals in pest management (Dube 2016) while the k -value has been extensively used as a measure of phytotoxic plant response (Pelinganga 2013; Mashela et al. 2015) and toxicity/efficacy of the allelochemicals to the pest (Dube et al. 2018). Usually, k -values start from zero and increase as discrete numbers when the sensitivity of the test organism to the allelochemical decreased (Liu et al. 2003).

Maerua angolensis and *Tabernaemontana elegans* plant extracts are used in traditional medicine for the treatment of various domestic animal and human ailments (Wink and Van Wyk 2008; Khosa 2013) and have shown potential in glasshouse, microplot and field trials in controlling *Meloidogyne* on tomato (Khosa 2013). Root-knot nematodes (*Meloidogyne*), on the other hand, are the most damaging nematode pests that parasitise crops (Jones et al. 2013). Information on the use of the CARD model to explain the relationship between the concentrations of the two plant extracts and plant-parasitic nematodes is not available. The objective of this study was to use the CARD model in the determination of *Meloidogyne incognita* sensitivity, DDG patterns and minimum inhibition concentrations of *M. angolensis* and *T. elegans* plant extracts on juvenile hatch and mortality.

Materials and methods

Collection of plant material and preparation of powdered leaf meals

Leaves of *M. angolensis* and *T. elegans* were collected from traditional healers in the Limpopo Province. The botanical origin of the leaf samples was identified and confirmed by a botanist from the South African National Biodiversity Institute (SANBI, Tshwane) and assigned voucher specimen numbers: *M. angolensis* 3112000/PRE099594-0 and *T. elegans* 6603000/PRE0812589-0. The leaf parts were chopped into 5 cm pieces and oven-dried for 4 days at 52°C prior to grinding in a Wiley mill and passing the material through a 1-mm-aperture sieve (Makkar 1999). Milled crude plant materials of 25 kg were bulk-stored in marked, air-tight glass containers at room temperature in the dark until used in the experiments.

Five hundred grams of crude leaf meal of *M. angolensis* and *T. elegans* were separately and independently left to soak in 1 L de-ionised water for 24 h

at room temperature. The extracts were filtered and the filtrates were freeze-dried to remove the water from the material.

Preparation of stock solutions

Ten stock solutions were prepared for each of the plant extract. Using an analytical balance, measures from 10 to 100 mg in increments of 10 mg were made from each extract and placed separately into sterile 1.5 mL Eppendorf tubes. This was followed by adding a solution of pluronic gel and de-ionised water to fill each tube to the 1 mL mark, after which the tubes were tightly closed. The extracts were brought into suspension in each tube by thoroughly shaking it for 45 min. The stock solutions were stored at -20°C in a freezer so that the stability of the extracts would not be affected.

Preparation of nematode inoculum

A population of *M. incognita* race 2, of which the identification was confirmed by sequence-characterised amplified regions-polymerase chain reaction (SCAR-PCR) (Zijlstra et al. 2000; Fourie et al. 2001), was obtained from the ARC-Grain Crops Institute, Potchefstroom, SA, and multiplied for a period of 2 months in glasshouse on susceptible tomato (*Solanum lycopersicum* L.) plants cv. 'Floradade'.

Dark-brown coloured egg masses of *M. incognita* were obtained from infected tomato plants, placed in 1% NaOCl solution and shaken for 30 s to surface-sterilise them and remove the gelatin matrix. The sterilised eggs were thoroughly rinsed in distilled water before used in J2 hatch bioassay.

Freshly-hatched J2 were obtained by transferring surface-sterilised eggs in Petri dishes containing 10 mL distilled water and placed in an incubator at 25 ± 2°C. J2 that hatched during the first 24 h were discarded with J2 that hatched during the subsequent 48 h used in the mortality bioassay (Dube and Mashela 2016).

Meloidogyne incognita second-stage juvenile mortality and hatch assays

In vitro trials were initiated under laboratory conditions at the Biosciences Unit at the Council for Scientific and Industrial Research (CSIR) in Pretoria, South Africa (approx. 25°44'47.96" S, 28°16'47.01" E).

Second-stage juvenile mortality assays: The concentrations of *M. angolensis* and *T. elegans* extracts were tested for nematode J2 mortality at ten different concentrations (0.1–1.0 mg mL⁻¹) each in 96-well test plates. This was achieved as follows: A suspension of 20 mL pure

pluronic gel and de-ionised water was put into 50 mL Eppendorf tubes. This was followed by adding a 10 mL suspension of J2 of *M. incognita* race 2 in water containing approximately 35,000 J2. The suspension was thoroughly shaken manually for 5 s. From this a 90 μL suspension containing 100 ± 20 J2 was transferred to the 96-well plate using a pipette. Ten microlitre extract of each concentration was subsequently put in each well plate in a specific direction and order. Thus, the final volume in each well was 100 μL . Treatments were arranged in a randomised complete block design (RCBD), replicated four times.

The ten treatment concentrations of each plant plus two controls of pure pluronic gel + deionised water solution and 1.8 mg mL⁻¹ salicylic acid dissolved in MeOH were assessed. Each well plate was sealed tightly with plastic wrapping to avoid any cross-well contamination and shaken for 10 min at 1000 revolutions per minute (RPM) on an Orbit 1000 (Labnet, Edison, USA) laboratory shaker. The number of living and dead J2 were determined prior to the onset of each experiment using an inverted compound microscope (Carl Zeiss, Jena Germany) set at 10X and 32X magnification. The plates were incubated at 22°C in the dark and the number of dead J2 per well plate was counted at 24, 48 and 72 h after the onset of the experiment. Two independent trials were conducted for each experiment.

Second-stage juvenile hatch assays: The exposure of eggs to different extracts and concentrations of *M. angolensis* and *T. elegans* were done with 100 ± 20 eggs in 96-well plates. Ten concentrations (0.1–1.0 mg mL⁻¹) of each plant extract plus two controls of pure pluronic gel + deionised water solution and 1.8 mg mL⁻¹ salicylic acid dissolved in MeOH were arranged in RCBD with four replications. Every treatment was randomly assigned to four rows (replicate blocks). After having sealed each well plate tightly with parafilm to avoid any cross-well contamination, each well plate was shaken for 10 min at 1000 RPM using an Orbit 1000 (Labnet, Edison, USA) laboratory shaker. The number of hatched J2 were determined prior to the onset of each experiment using an inverted compound microscope (Carl Zeiss, Jena Germany) set at 10x and 32x magnification. The plates were incubated at 22°C in the dark and the number of hatched J2 per well plate were counted at 7, 14 and 21 days after the onset of experiment. Two independent trials were conducted for each plant extract.

Statistical analysis

Cumulative J2 percentage hatch and mortality were made per treatment after each incubation period. The standardised residuals of each variable were tested for

deviations from normality using Shapiro-Wilk's test. Data were transformed using arc-sine (percentage)^{1/2} prior to analysis of variance using SAS/STAT statistical software (SAS Institute 2008). Fisher's protected t-LSD (least significant difference) was calculated at 5% level of significance to compare means of significant effects (Snedecor and Cochran 1980). Significant mean exposure period values were subjected to the CARD model (Liu et al. 2003) to generate the J2 hatch and mortality curves using the quadratic equation $Y = b_2x^2 + b_1x + c$, where Y = J2 hatch inhibition or mortality mean value and x = plant extract concentration mean value. Additionally, the CARD-generated biological indices, namely threshold stimulation (D_m), saturation point (R_h), 0% inhibition concentration (D_0), 50% inhibition concentration (D_{50}), 100% inhibition concentration (D_{100}), sensitivity index (k) and coefficient of determination (R^2) (Liu et al. 2003), were summarised. Unless otherwise stated, only treatments that were significant at 5% probability level were discussed.

Results

Maerua angolensis and Tabernaemontana elegans plant extracts on juvenile hatch

The CARD model quantified concentration ranges of the two plant extracts that could stimulate ($D_m - R_h$), saturate ($R_h - D_0$) and inhibit ($D_0 - D_{100}$), J2 hatch (Table 1). The CARD-generated DDG patterns demonstrated an increase in juvenile hatch inhibition with the increase in concentration of the plant extracts (Figure 1). The DDG patterns were explained by 74%, 93% and 97% of the derived models at 7-, 14- and 21-day exposure periods, respectively, for *M. angolensis* and 87%, 98% and 99%, at the same exposure periods for *T. elegans* (Table 1). The sensitivity of J2 hatch to increasing concentrations

Table 1. CARD-generated biological indices of *Meloidogyne incognita* second-stage juvenile hatch to increasing concentrations of *Maerua angolensis* and *Tabernaemontana elegans*.

Biological index	<i>Maerua angolensis</i>			<i>Tabernaemontana elegans</i>		
	Exposure period (days)			Exposure period (days)		
	7	14	21	7	14	21
Threshold stimulation (D_m)	–	0.265	0.329	0.122	0.169	0.173
Saturation point (R_h)	0.636	1.001	1.136	0.584	0.973	1.001
0% inhibition (D_0)	0.636	1.266	1.465	0.706	1.142	1.174
50% inhibition (D_{50})	0.636	1.266	1.465	0.706	1.142	1.174
100% inhibition (D_{100})	0.636	1.266	1.465	0.706	1.142	1.174
R^2	0.74	0.93	0.97	0.87	0.98	0.99
Sensitivity index (k)	20	20	20	20	17	19
Average sensitivity ranking (k)		20			19	




Figure 1. Density-dependent growth patterns of *Meloidogyne incognita* juvenile hatch at 7, 14 and 21 days of incubation in *Tabernaemontana elegans* (A) and *Maerua angolensis* (B).

of *M. angolensis* and *T. elegans* was moderate at an average sensitivity ranking (k) of 20 and 19 units, respectively (Table 1). The minimum juvenile hatch inhibition concentration for the two extracts were comparable at 0.06 and 0.07 mg mL⁻¹ for *M. angolensis* and *T. elegans*, respectively (Table 2).

Maerua angolensis and Tabernaemontana elegans plant extracts on juvenile mortality

The effects of the two plant extracts on *M. incognita* juvenile mortality were similar to those of juvenile hatch in terms of the CARD model quantified concentration ranges that could stimulate ($D_m - R_h$), saturate ($R_h - D_0$) and inhibit ($D_0 - D_{100}$), J2 mortality (Table 3) and DDG patterns (Figure 2). The DDG patterns were explained by 92, 85% and 96% of the derived models at 24-, 48- and 72-h exposure periods, respectively for *M. angolensis* and 89%, 94% and 70% at the same exposure periods for *T. elegans* (Table 3). Second-stage juvenile mortality was highly sensitive to *M. angolensis* with average k -value of 2 units, whereas the sensitivity to *T. elegans* was moderate at k -value of 15 units (Table 3). The minimum juvenile

mortality concentration for the two extracts were very low at 0.13 and 0.94 $\mu\text{g mL}^{-1}$ for *M. angolensis* and *T. elegans*, respectively (Table 4).

Discussion

Maerua angolensis and Tabernaemontana elegans plant extracts on Meloidogyne incognita juvenile hatch

The nematode suppressive nature of the two plant extracts used in traditional medicine in this study is similar to reports made previously on other plants (Ntalli and Caboni 2012). Several plant extracts used in traditional medicine have demonstrated a measure of bioactivity on nematode J2 hatch (Ibrahim et al. 2006; Javed et al. 2008; Kalaiselvam and Devaraj 2011). The most studied plant species include, garlic (*Allium sativum* L.), chrysanthemum (*Chrysanthemum coronarium* L.), fennel (*Foeniculum vulgare* Mill) (Ibrahim et al. 2006), mugwort (*Artemisia vulgaris* L.) (Costa et al. 2003), *Azadirachta indica* (Javed et al. 2008), *Inula viscosa* (Oka et al. 2001), white cedar (*Melia azedarach* L.), elderberry (*Sambucus nigra* L.) (Akyazi 2014) and *Tagetes* sp. (Kalaiselvam and

Table 2. *Maerua angolensis* and *Tabernaemontana elegans* minimum *Meloidogyne incognita* juvenile hatch inhibition concentration at three exposure periods.

Exposure period (days)	<i>Maerua angolensis</i>		<i>Tabernaemontana elegans</i>	
	Model	x (mg mL ⁻¹) ^a	Model	x (mg mL ⁻¹)
7	$y = 0.30 + 11.16x - 48.94x^2$	0.11	$y = 0.71 + 21.09x - 190.37x^2$	0.06
14	$y = 0.52 + 27.16x - 184.20x^2$	0.07	$y = 0.71 + 27.79x - 198.40x^2$	0.07
21	$y = 0.52 + 29.09x - 186.29x^2$	0.08	$y = 0.77 + 30.24x - 228.44x^2$	0.07
Mean		0.08		0.07

^a $x = -b_1/2b_2$, where $y = b_2x^2 + b_1x + c$.

Table 3. Biological indices of *Meloidogyne incognita* second-stage juvenile mortality to increasing concentrations of *Maerua angolensis* and *Tabernaemontana elegans*.

Biological index	<i>Maerua angolensis</i>			<i>Tabernaemontana elegans</i>		
	Exposure period (h)			Exposure period (h)		
	24	48	72	24	48	72
Threshold stimulation (D_m)	1.008	0.626	–	–	0.888	0.265
Saturation point (R_p)	0.992	1.124	2.433	0.921	0.886	0.918
0% inhibition (D_0)	15.602	1.750	2.433	0.921	1.774	1.183
50% inhibition (D_{50})	15.821	1.750	2.433	0.921	1.774	1.183
100% inhibition (D_{100})	16.00	1.750	2.433	0.921	1.774	1.183
R^2	0.92	0.85	0.96	0.89	0.94	0.70
Sensitivity index (k)	3	0	3	17	9	20
Average sensitivity ranking (k)		2			15	

Devaraj 2011). In most of these plants, DDG responses have been reported with majority of these observations showing an inverse relationship between J2 hatch and the increasing concentrations (Javed et al. 2008; Kalaiselvam and Devaraj 2011; Akyazi 2014).

Mashela et al. (2015) reported that generally, biological systems respond to extrinsic and intrinsic factors in DDG patterns, characterised by three growth phases, namely, stimulation, saturation (neutral) and inhibition in support of the theory postulated by Liu et al. (2003). The CARD model was developed to quantify these DDG response patterns of biological entities to increasing concentration of allelochemicals (Liu et al. 2003). The stages of DDG patterns were observed on *M. incognita* J2 hatch, mobility and mortality exposed to increasing concentrations of fermented crude extracts of *Cucumis myriocarpus* and *Cucumis africanus* (Dube and

Mashela 2017; Dube et al. 2018). All three stages of DDG patterns were observed in the current study when *M. incognita* J2 were exposed to *M. angolensis* and *T. elegans* plant extracts (Table 1 and Figure 1) supporting previous documentations (Liu et al. 2003; Mashela et al. 2015). The observations in the current study provided evidence for the potential use of the CARD model as previously reported (Mashela et al. 2015).

The higher the sensitivity (k) value of an organism, the higher the tolerance of that organism to allelochemicals, *vice versa* (Liu et al. 2003). In the current trials, J2 hatch was moderately sensitive to both *M. angolensis* and *T. elegans* plant extracts (Table 1). In plant-parasitic nematodes, successful J2 hatch and other behaviours depend on the ability of the chemical concentrations in soil solutions to enter and be detected by chemoreceptors in developing nematode inside eggs (McSorley 2003). The moderate sensitivity effects of the two plant extracts could be explained in terms of the role played by the layers of the eggs that could have provided some kind of restrictions to the movement of chemicals hence affecting the concentrations to which the nematode inside the egg is exposed to (McSorley 2003).

In the current study, minimum J2 hatch inhibition of *M. angolensis* and *T. elegans* plant extracts were very low at all exposure periods (Table 2) compared to previous studies (Dube and Mashela 2016). The findings contradict the previous report that low minimum inhibition concentrations are associated with high extract sensitivities (Dube 2016). In the current study, minimum J2 hatch inhibition concentrations were low, whereas the sensitivity to the two plant extracts was moderate.




Figure 2. Density-dependent growth patterns of *Meloidogyne incognita* juvenile mortality at 24, 48 and 72 h of incubation in *Tabernaemontana elegans* (A) and *Maerua angolensis* (B).

Table 4. *Maerua angolensis* and *Tabernaemontana elegans* minimum *Meloidogyne incognita* juvenile mortality concentration at three exposure periods.

Exposure period (h)	<i>Maerua angolensis</i>		<i>Tabernaemontana elegans</i>	
	Model	x (mg mL ⁻¹) ^a	Model	x (mg mL ⁻¹)
24	$y = 0.01 + 4.67x - 5.50x^2$	0.42	$y = 0.03 + 13.23x - 47.51x^2$	0.14
48	$y = 0.16 + 3.59x - 2.87x^2$	0.63	$y = 0.12 + 9.52x - 25.59x^2$	0.19
72	$y = 0.52 + 2.75x - 0.78x^2$	1.76	$y = 0.24 + 24.91x - 168.95x^2$	0.07
Mean		0.94		0.13

^a $x = -b_1/2b_2$, where $y = b_2x^2 + b_1x + c$.

Maerua angolensis and Tabernaemontana elegans plant extracts on Meloidogyne incognita juvenile mortality

The bioactive properties of *M. angolensis* and *T. elegans* plant extracts used in traditional medicine is similar to previous observations of *in vitro* studies of essential oil from true myrtle (*Myrtus communis* L.) that exhibited high mortalities of *M. incognita* (Archana and Prasad 2014). Various other plant extracts from *Datura stramonium*, *A. indica*, *Moringa* species, *Artemisia vulgaris* and *Allium sativum* all displayed lethal properties to *Meloidogyne* species (Ibrahim et al. 2006; Claudius-Cole et al. 2010; Nelaballe and Mukkara 2013).

The J2 mortality DDG patterns observed in this study (Figure 2) agreed with the previously postulated hypothesis on the response of organisms to increasing concentrations of various allelochemicals (Liu et al. 2003). Pure cucurbitacin A and B also exhibited all stages of DDG patterns on *M. incognita* J2 mortality when the CARD model was used (Dube and Mashela 2016), findings similar to current study.

The minimum lethal concentration due to *M. angolensis* and *T. elegans* observed in this study were very low (Table 4) compared to previous studies (Ardakani et al. 2013). Essential oils at concentrations as high as 250 mg mL⁻¹ could not cause mortality of *M. incognita* J2 (Ardakani et al. 2013) compared to averages of 0.13 and 0.94 mg mL⁻¹ observed in the current study. For pure isothiocyanate, aldehydes and ketones, *M. incognita* J2 mortality were observed at very low concentrations of 5, 15 and 13 µL L⁻¹, respectively (Ntalli and Caboni 2012) compared to crude plant extracts in the current study.

Even though the *M. incognita* J2 hatch sensitivities to *M. angolensis* and *T. elegans* were both moderate and comparable, the sensitivity of J2 mortality to the two plant extracts differed (Table 3). *Meloidogyne incognita* J2 mortalities were more sensitive to *M. angolensis* than to *T. elegans*.

The difference in effects of the same plant extracts on *M. incognita* eggs and J2 can be explained in terms of the role played by the layers of the eggs that could have

provided restrictions to the movement of chemicals into the egg where a developing nematode is found and the properties of the active substances in the extracts to penetrate these layers (Dube 2016). This heightened sensitivity of J2 to plant extracts when compared to eggs has been reported in a number of other studies (Javed et al. 2008; Akyazi 2014).

In conclusion, CARD model was able to generate the DDG patterns, sensitivity values and minimum inhibition concentrations of *M. angolensis* and *T. elegans*. *Meloidogyne incognita* J2 hatch inhibition and mortality to *M. angolensis* and *T. elegans* observed in this study demonstrated the nematicidal properties of the two products. The low minimum J2 hatch, lethal concentrations, and moderate sensitivity values observed in this study provides evidence of potentially high potency of the *M. angolensis* and *T. elegans* when compared to other plant-based compound.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This study was financially supported by the Department of Science and Technology Republic of South Africa and the National Research Foundation Republic of South Africa.

Notes on contributors

M. C. Khosa (PhD) Researcher at Agricultural Research Council-Tropical and Subtropical Crops, Nelspruit, South Africa.

Z. P. Dube (PhD) Lecturer in the School of Biology and Environmental Sciences, University of Mpumalanga, Mbombela, Republic of South Africa.

M. Tselanyane (PhD) Senior Researcher at Biosciences Unit, Council of Scientific and Industrial Research, Pretoria, South Africa.

G. Fouche (PhD) Professor and Chief Research Scientist at Department of Chemistry, University of Pretoria, Pretoria, 0001, Republic of South Africa.

D. De Waele (PhD) Emeritus Full Professor in the Department of Biosystems, Faculty of Bioscience Engineering, University of Leuven (KU Leuven), Leuven, Belgium.

M. S. Daneel (PhD) Senior Researcher at Agricultural Research Council-Tropical and Subtropical Crops, Nelspruit, South Africa.

References

- Akyazi F. 2014. Effect of some plant methanol extracts on egg hatching and juvenile mortality of root-knot nematode *Meloidogyne incognita*. *Am J Exp Agric*. 4:1471–1479.
- Archana US, Prasad D. 2014. Management of plant-parasitic nematodes by the use of botanicals. A review. *J Plant Physiol Pathol*. 2:9–16.
- Ardakani AS, Hosyninej SA, Pourshirz A. 2013. Killing effects of *Myrtus communis* L. essential oil on *Meloidogyne incognita*. *Int J Agri Crop Sci*. 5:806–810.
- Claudius-Cole AO, Aminu AE, Fawole B. 2010. Evaluation of plant extracts in the management of root-knot nematode *Meloidogyne incognita* on cowpea (*Vigna unguiculata* (L) Walp). *Mycopath*. 8:53–60.
- Costa SSR, Santos MSNA, Ryan MF. 2003. Effect of *Artemisia vulgaris* rhizome extracts on hatching, mortality, and plant infectivity of *Meloidogyne megadora*. *J Nematol*. 35:437–442.
- Donatelli M, Magarey RD, Bregaglio S, Willocquet L, Whish JPM, Savary S. 2017. Modelling the impacts of pests and diseases on agricultural systems. *Agric Syst*. 155:213–224.
- Dube ZP. 2016. Nema-AL and Nema-BL phytonematicides: bioactivities in *Meloidogyne incognita*, tomato crop, soil type and organic matter [PhD thesis]. Polokwane: University of Limpopo.
- Dube ZP, Mashela PW. 2016. Response of *Meloidogyne incognita* egress and overall sensitivity to active ingredients of Nema-AL and Nema-BL phytonematicides. *Transylv Rev*. 24(7):954–960.
- Dube ZP, Mashela PW. 2017. Developing cucurbitacin A minimum inhibition concentration for *Meloidogyne incognita* using a computer-based model. *Transylv Rev*. 25(19):5939–5943.
- Dube ZP, Mashela PW, De Waele D. 2018. Comparison of the activity of pure cucurbitacins and cucurbitacin-containing crude-extracted phytonematicides on the hatching, motility and viability of the infective second-stage juveniles of *Meloidogyne incognita*. *S Afr J Plant Soil*. 36(1):29–32.
- Fisker EN, Bak J, Niassy A. 2007. A simulation model to evaluate control strategies for the grasshopper, *Oedaleus senegalensis* in West Africa. *Crop Prot*. 26:592–601.
- Fourie H, Zijlstra C, McDonald AH. 2001. Identification of root-knot nematodes species occurring in SA using the SCAR-PCR technique. *Nematology*. 3:675–680.
- Gutsche V, Rossberg D. 1997. SYNOPSIS 1.1: a model to assess and to compare the environmental risk potential of active ingredients in plant protection products. *Agric Ecosyst Environ*. 64:181–188.
- Herbert DA, Mack TP, Backman PA, Rodriguez-Kabana R. 1992. Validation of a model for estimating leaf-feeding by insects in soybean. *Crop Prot*. 11:27–34.
- Ibrahim SK, Traboulsi AF, El-Haj S. 2006. Effect of essential oils and plant extracts on hatching, migration and mortality of *Meloidogyne incognita*. *Phytopathol Mediterr*. 45:238–246.
- Javed N, Gowen SR, El-Hassan SA, Inam-Ul-Haq M, Shahina F, Pembroke B. 2008. Efficacy of neem (*Azadirachta indica*) formulations on biology of root-knot nematodes (*Meloidogyne javanica*) on tomato. *Crop Prot*. 27:36–43.
- Jones JT, Haegeman A, Danchin EGJ, Gaur HS, Helder J, Jones MGK, Kikuchi T, Manzanilla-Lopez R, Palomares-Rius JE, Wesemael WML, et al. 2013. Top 10 plant-parasitic nematodes in molecular plant pathology. *Mol Plant Pathol*. 14:946–961.
- Kalaiselvam I, Devaraj A. 2011. Effect of root exudates of *Tagetes* sp. on egg hatching behaviour of *Meloidogyne incognita*. *Int Res J Pharm*. 2:93–96.
- Khosa MC. 2013. An investigation into the potential of crude and partially separated material of selected non-crop plant species as control agents of root-knot nematodes (*Meloidogyne incognita*) in tomato [PhD thesis]. Potchefstroom: North-West University.
- Liu DL, An M, Johnson IR, Lovett JV. 2003. Mathematical modeling of allelopathy. III. A model for curve-fitting allelochemical dose responses. *Nonlinearity Biol Toxicol Med*. 1:37–50.
- Mafeo TP. 2012. Responses of economically important crops to crude extracts of cucumis fruit when used as pre-emergent bio-nematicide [PhD thesis]. Sovenga: University of Limpopo.
- Makkar HPS. 1999. *Quantification of tannins in tree foliage: a laboratory manual for the FAO/IAEA co-ordinated research project on use of nuclear and related techniques to develop simple tannin assay for predicting and improving the safety and efficiency of feeding ruminants on the Tanniniferous tree foliage*. Vienna: IAEA Working Document. p. 29.
- Mashela PW, Dube ZP, Pofu KM. 2015. Phytotoxicity of soil-amended phytonematicides and related inconsistent results on nematodes suppression. In: Meghvanji MK, Vormaa A, editor. *Organic amendments and soil suppressiveness*. Cham: Springer International Publishers; p. 147–173.
- McSorley R. 2003. Adaptations of nematodes to environmental extremes. *Fla Entomol*. 86:138–142.
- Nayak P, Mukherjee AK, Pandit E, Pradhan SK. 2018. Application of statistical tools for data analysis and interpretation in rice plant pathology. *Rice Sci*. 25(1):1–8.
- Nelaballe VK, Mukkara LD. 2013. A preliminary study on the nematocidal effect of some local flora on *Meloidogyne incognita* chitwood infesting mulberry. *Int J Chem Environ Biol Sci*. 1:475–477.
- Ntalli NG, Caboni P. 2012. Botanical nematicides. A review. *J Agric Food Chem*. 60:9929–9940.
- Oka Y, Ben-Daniel B, Cohen Y. 2001. Nematicidal activity of powder and extracts of *Inula viscose*. *Nematologica*. 3:735–742.
- Pelinganga OM. 2013. Developing bio-nematicides using indigenous cucumis africanus and cucumis myriocarpus fruits for tomato production system [PhD thesis]. Polokwane: University of Limpopo.
- SAS Institute. 2008. *SAS/STAT 9.2 qualification tools user's guide*. Cary, NC: SAS Institute.
- Snedecor GW, Cochran WG. 1980. *Statistical methods*. 7th ed. Ames: Iowa State University Press; p. 507.
- Walczak F. 2011. Applying a computer program, based on a developed mathematical model, to protect cereal crops from leaf beetles (*Oulema* spp.). *Arch Phytopathol Plant Protect*. 44(13):1238–1247.
- Wink M, Van Wyk BE. 2008. *Mind-altering and poisonous plant of the world*. Pretoria: Briza; p. 268.
- Zijlstra C, Donkers-Venne DTHM, Fagette M. 2000. Identification of *Meloidogyne incognita*, *M. javanica* and *M. arenaria* using sequence characterized amplified region (SCAR) based PCR. *Nematology*. 2:847–853.